

Original Research Article

STUDY OF SPECTRUM OF HEMATOLOGICAL DISORDERS IN BONE MARROW ASPIRATION AND TREPHINE BIOPSY IN TERTIARY CARE CENTRE

Ajay Kumar Gupta¹, Sumit Kumar², Shivangi shukla³, Kajal Godara⁴, Pooja Purohit⁵, Naresh N rai⁶

¹Assistant Professor, Department of Pathology, Jaipur National University Institute for Medical Sciences & Research Centre, Jaipur, Rajasthan, India.
 ²Assistant Professor, Department of Pathology, Jaipur National University Institute for Medical Sciences & Research Centre, Jaipur, Rajasthan, India.
 ³PG Junior Resident, Department of Pathology, Jaipur National University Institute for Medical Sciences & Research Centre, Jaipur, Rajasthan, India.
 ⁴PG Junior Resident, Department of Pathology, Jaipur National University Institute for Medical Sciences & Research Centre, Jaipur, Rajasthan, India.
 ⁴PG Junior Resident, Department of Pathology, Jaipur National University Institute for Medical Sciences & Research Centre, Jaipur, Rajasthan, India.
 ⁵Senior Resident, Department of Pathology, Jaipur National University Institute for Medical Sciences & Research Centre, Jaipur, Rajasthan, India.
 ⁶Professor & Head, Department of Pathology, Jaipur National University Institute for Medical Sciences & Research Centre, Jaipur, Rajasthan, India.

 Received
 : 04/03/2025

 Received in revised form : 30/04/2025
 Accepted

 Accepted
 : 17/05/2025

Corresponding Author:

Dr. Ajay Kumar Gupta, Assistant Professor, Department of Pathology, Jaipur National University Institute for Medical Sciences & Research Centre, Jaipur, Rajasthan. Email: dr.gupta.ajay17@gmail.com

DOI: 10.70034/ijmedph.2025.2.360

Source of Support: Nil, Conflict of Interest: None declared

Int J Med Pub Health 2025; 15 (2); 2009-2012

ABSTRACT

Background: Blood is an opaque fluid consists of plasma, a pale-yellow, coagulable fluid, in which different types of blood cells are suspended. These cells include erythrocytes, granulocytes, lymphocytes, monocytes, and platelets. There are numerous blood and blood related disorders which can be diagnosed by detailed clinical examination and laboratory investigations such as complete blood count, peripheral blood smear along with biochemical analysis and radiological imaging. When peripheral smear is not informative, further bone marrow examination and higher investigations such as cytochemistry, immunophenotyping, cytogenetic analysis and molecular genetic studies can aid for arriving at a definitive diagnosis.

Materials and Methods: A prospective, descriptional study was conducted in the department of pathology, jaipur national university institute for medical science & research centre, jaipur, rajasthan between the period of January 2024 to Jan 2025. A total number of 100 cases were evaluated who underwent both bone marrow aspiration and biopsy. The patients with negative bone marrow aspiration and unsatisfactory bone marrow biopsy were excluded.

Results: Our study enlists the spectrum of lesions encountered in bone marrow aspiration and biopsy findings in various haematological and non-haematological disorders. The diagnostic efficacy of bone marrow trephine biopsy over the aspiration smear was evaluated. A total of 100 cases were studied. Among the 100 cases studied, 94% were haematological disorders followed by 4% non-haematological disorders and 2% were normal bone marrow. The age of the patients ranged from 9 to 85 years, with a mean age of 50.22 years. The most common age group of presentation in our study was 50 to 59 years, which accounted for 23% of the cases. Most of the patients were males accounting for 63% (63 cases), with male to female ratio of 1.7:1.

Conclusion: Bone marrow examination is helpful in arriving at final diagnosis of various haematological and nonhaematological lesions. It consists of bone marrow aspiration and bone marrow biopsy which can be done in the single sitting. The bone marrow biopsy remains the gold standard for the diagnosis in negative aspiration. Simultaneous evaluation of bone marrow aspiration and biopsy show good correlation and increases the efficacy of diagnosis.

Keyword: Bone Marrow Aspiration(BMA), Trephine Biospy.

INTRODUCTION

Blood is an opaque fluid consists of plasma, a paleyellow, coagulable fluid, in which different types of blood cells are suspended. These cells include erythrocytes, granulocytes, lymphocytes, monocytes, and platelets.^[1] There are numerous blood and blood related disorders which can be diagnosed by detailed clinical examination and laboratory investigations such as complete blood count. peripheral blood smear along with biochemical analysis and radiological imaging. When peripheral blood smear is not informative, further bone marrow examination and higher such cytochemistry, investigations as immunophenotyping, cytogenetic analysis and molecular genetic studies can aid for arriving at a definitive diagnosis.^[2] This study encompasses the spectrum of lesions encountered in bone marrow aspiration and trephine bone marrow biopsy findings in various haematological and nonhaematological disorders. The diagnostic efficacy of bone marrow trephine biopsy over the aspiration smear was evaluated.

MATERIALS AND METHODS

A prospective, descriptional study was conducted in the department of pathology, jaipur national university institute for medical science & research centre, jaipur, rajasthan between the period of January 2024 to Jan 2025. A total number of 100 cases were evaluated who underwent both bone marrow aspiration and biopsy. The patients with negative bone marrow aspiration and unsatisfactory bone marrow biopsy were excluded. The clinical data was recorded including physical examination, relevant haematological, biochemical and radiological investigations. The posterior superior iliac spine was the most suitable site used for both aspiration and biopsy. The procedure was done under local anesthesia with 2% injection lignocaine. Bone marrow aspiration slides were stained with Leishman stain and bone marrow biopsy bits were kept for decalcification for 24 hours and then processed and stained with haematoxylin and eosin stains. Special stains were performed when ever required. Data was analyzed by using statistical tools: SPSS software version 22.0 and CHI-square test to evaluate sensitivity, specificity, positive predictive value and negative predictive value. Comparision of quantitative variables were done using t-test. Significance were considered if p value is less than 0.05.

RESULTS

Our study enlists the spectrum of lesions encountered in bone marrow aspiration and biopsy findings in various haematological and nonhaematological disorders. The diagnostic efficacy of bone marrow trephine biopsy over the aspiration smear was evaluated. A total of 100 cases were studied. Among the 100 cases studied, 94% were haematological disorders followed by 4% nonhaematological disorders and 2% were normal bone marrow. Table 1, shows distribution of various haematological disorders and Table 2, shows various non- haematological lesions. The age of the patients ranged from 9 to 85 years, with a mean age of 50.22 years. The most common age group of presentation in our study was 50 to 59 years, which accounted for 23% of the cases. Most of the patients were males accounting for 63% (63 cases), with male to female ratio of 1.7:1. The p- value of both bone marrow aspiration and bone marrow biopsy in relation to sex was found to be statistically significant < 0.05.

2010

Table 1: Distribution of various haematological disorders				
S. No	Haematological disorders	Number	Percentage (%)	
1	Anaemia	31	32.97	
2	Plasma cell disorder	16	17.02	
3	Myeloproliferative neoplasms	15	15.95	
4	Lymphoproliferative disorder	06	6.38	
5	Acute myeloid leukaemia	12	12.76	
6	Acute lymphoid leukaemia	04	4.25	
7	Myelodysplastic syndrome	02	2.12	
8	Myelodysplastic syndrome/myeloproliferative neoplasms	01	1.06	
9	Immune thrombocytopenic purpura	06	6.38	
10	Acute myeloid leukemia with myelodysplastic syndrome	01	1.06	
	related changes			
Total		94	100	

Table 2: Various non-haematological lesions				
S. No	Various non-haematological lesions	Number	Percentage (%)	
1	Granulomatous lesion	2	50	
2	Metastatic solid tumour deposit	1	25	
3	Infection	1	25	
Total		4	100	



Figure 1: (A) Iron Deficiency Anaemia: Aspiration showing micro-normoblast maturation (100x), inset showing micro-normoblasts; (B) Megaloblastic Anaemia: Aspiration showing megaloblasts displaying sieve-like chromatin of the (40x), inset showing megaloblasts; (C) Multiple Myeloma: Aspiration showing sheets of plasma cells with few binucleated forms [arrow]. (100x); (D) Metastatic Deposits: Biopsy showing pleomorphic tumour cells (H&E 40x)

DISCUSSION

The clinical diagnosis at times may need complete blood count, peripheral blood smear for arriving at a final diagnosis. When peripheral blood smear is not informative, further bone marrow examination and higher investigations may be essential for a definitive diagnosis.

It is a useful investigative tool to diagnose many haematological and non-haematological disorders and thereby, enabling to formulate an effective treatment, staging and management of malignant lesions. A total number of hundred (100) cases were studied. The majority of cases encountered were haematological disorders accounting for 94% followed by 4% of non-haematological disorders and 2% were normal bone marrow. These findings were in concordance with various studies by Mahajan, et al,^[3] Khatik, et al,^[4] and Goyal, et al.^[5] The age range was between 9 to 85 years, with the mean age being 50.22 years. The age distribution was comparable to the studies conducted by Panigrahi, et al,^[6] Khan, et al,^[7] Tyagi, et al,^[8] and Atchyuta, et al.^[9]

The number of male patients in the present study were 63% and the female patients were 37% with a male to female ratio of 1.7:1. This was in

concordance with the studies done by Goyal, et al,^[5] Panigrahi, et al,^[6] Khan, et al,^[7] Atchyuta, et al,^[9] and Birare, et al.^[10]

Anaemia was seen in 32.97% cases wherein, bone marrow aspiration was satisfactory in 95.83%, whereas, biopsy was satisfactory in 100% cases. However, aspiration was dry tap in 4.17% cases whereas biopsy was cellular. The most common anaemia encountered was megaloblastic anaemia, which was comparable with the studies by Khatik, et al,^[4] Atchyuta, et al,^[9] Birare, et al,^[10] Bashir, et al,^[11] and Mehra, et al.^[12]

The second commonest anaemia encountered was aplastic anaemia, which was in concordance with the studies by Khatik, et al,^[4] Atchyuta, et al,^[9] and Mehra, et al.^[12] The incidence of nutritional anaemia noted in the present study was higher, probably because of higher prevalence of nutritional deficiencies. Plasma cell disorders were encountered in 17.02% of cases.

The most common disorder observed was Multiple myeloma (MM) with 86.35% of cases, followed by 4.55% of cases each of Plasma cell leukaemia (PCL), Waldenstrom macroglobunemia (WM) and Monoclonal gammopathy of undetermined significance (MGUS), which was in concordance with the studies by Atchyuta, et al,^[9] Birare, et al,^[10] and Vijaymohanan, et al,^[13] (Figure 1).

The basic panel markers used in myeloma are CD19, CD56, CD38, CD138, CD81, CD27, CD45, cytoplasmic kappa and lambda. Among 15.95% of cases of MPNs, bone marrow aspiration was satisfactory in 68.75% of cases, and rest 31.25% were dry tap. The cause of dry tap in MPNs was probably due to increased bone marrow fibrosis or cellularity.

Biopsy was satisfactory and cellular in all the cases. Bone marrow biopsy not only helps in differentiation of MPN, but also to assess cellularity, histo-topographic cell distribution, morphology of blasts as well as megakaryocytes and degree of marrow fibrosis. The most common MPN encountered in the present study was Chronic Myeloid Leukaemia (CML) accounting for 68.75% of cases, followed by Polycythaemia Vera (PV) in 12.50% of cases, 6.25% cases each of Essential Thrombocythemia (ET), Myelofibrosis (MF) and Myeloproliferative Neoplasm-unclassified (MPN-U).

In the studies conducted by Khatik, et al,^[4] and Atchyuta, et al,^[9] the most common disorder encountered was CML as in the present study. Biopsy gives additional information about pattern of involvement and prognosis. The detection of BM involvement virtually always advances lymphoma patients to Stage IV disease. Lymphoproliferative disorders were encountered in 6.38%. chronic lymphocytic leukaemia/ lymphoma was encountered in 50% cases, followed by Non-Hodgkin's lymphoma 33.33% cases of, and 16.67% of cases were Hodgkin's lymphoma infiltrating marrow.

The basic panel markers used in NHL, and CLL are CD19, CD45, FMC7, CD23, CD200, CD5, CD20, CD22, CD10, CD34, CD7, CD5, CD4, CD3, CD8, ZAP70, surface kappa, surface lambda, CD103, CD123, CD11c, TCR alpha-gamma/gamma-delta, CD26, and CD25. The prognostic markers used in CLL are CD38 and ZAP70.^[14,15]

Non-haematological disorders were encountered in 4% of cases. The most common lesions encountered were granulomatous lesions in 50% of cases followed by 25% cases of metastatic solid tumour deposits and 25% cases of infection. In our study, most common non-haematological disorder encountered was granulomatous lesion followed by metastatic solid tumours which was in concordance with the studies conducted by Khatik, et al,^[4] and Vijaymohanan, et al.^[13]

The sensitivity of bone marrow aspiration and biopsy was 92% in our study which was comparable with the studies carried out by Khatik, et al,^[4] Lukas B, et al,^[16] and Chauhan, et al.^[17] The diagnostic accuracy of bone marrow aspiration noted was 92% and biopsy was 100%.

The discrepancy rate between bone marrow aspiration and biopsy was 8%. The diagnostic accuracy of bone marrow aspiration was found to be lower as few cases (8 cases) showed unsatisfactory marrow aspirate/dry tap due to faulty technique.

The p-value of bone marrow aspiration and biopsy in relation to diagnosis is significant statistical, the p-value < 0.001 by chi-square method which was comparable with Goyal, et al,^[5] Bashir, et al,^[11] and Chauhan, et al.^[17]

CONCLUSION

Bone marrow examination is helpful in arriving at final diagnosis of various haematological and nonhaematological lesions. It consists of bone marrow aspiration and bone marrow biopsy which can be done in the single sitting. In bone marrow aspiration morphology of the cells are well preserved, whereas cell distribution, tumour infiltration, granuloma cells and storage cells are better appreciated on bone marrow biopsy. The bone marrow biopsy remains the gold standard for the diagnosis in negative aspiration. Simultaneous evaluation of bone marrow aspiration and biopsy show good correlation and increases the efficacy of diagnosis.

REFERENCES

- Wickramasinghe SN, Erber WN (2011) Normal blood cells. In: Porwit, Mc Cullough J, Erber WN, Blood and bone marrow pathology. (2nd edn), Edinburgh: Elsevier Limited, 3-18.
- Lee SH, Erber WN, Porwit A, Tomonaga M, Peterson LC (2008) ICSH guidelines for the standardization of bone marrow specimens and reports. Int J Lab Haematol 30: 349-364
- Mahajan V, Kaushal V, Thakur S, Kaushik R (2013) A comparative study of bone marrow aspiration and bone marrow biopsy in haematological and non-haematological disorders - An institutional experience. JIACM 14: 133-135.
- Khatik DK, Mishra D, Choudhary M (2017) A relevance study of bone marrow aspiration and bone marrow biopsy in haematological and non-haematological disorders. JMSCR 5: 20900-20908.
- Goyal N, Kundal RK, Singh H, Raman, Ashima (2018) Comparative study of bone marrow aspiration and bone marrow trephine biopsy in haematological and nonhaematological disorders. Ann Int Med Den Res 4: 65-69.
- Panigrahi P, Behera PK, Tripathy KP (2017) Bone marrow involvement in non-haematological disorders. An Institutional Experience 16: 51-53.
- Khan SP, Geelani S, Akhter S, Bhat S, Hussain S, et al. (2018) Bone marrow aspiration in haematological disorders: Study at a tertiary care centre. Int J Res Med Sci 6: 2361-2364.
- Tyagi R, Singh A, Garg B, Sood N (2018) Beware of bone marrow: Incidental detection and primary diagnosis of solid tumours in bone marrow aspiration and biopsies; A study of 22 cases. Iran J Pathol 13: 78-84.
- Atchyuta M, Premalatha P, Renuka IV, Krishnamashary PAV, Tejeswini V (2019) Spectrum of hematological diseases diagnosed by bone marrow examination in a tertiary care hospital. Ind J Pathol Oncol 6: 185-189.
- Birare SD, Jain N, Abhange RS (2019) A study of hematological indices and bone marrow morphology in patients presenting with anemia. Indian J Pathol Oncol 6: 266-274.
- Bashir N, Musharaf B, Reshi R, Jeelani T, Rafiq D, et al. (2018) Bone marrow profile in haematological disorders: An experience from a tertiary care centre. Int J Adv Med 5: 608-613.
- Mehra V, Rani B, Singh J, Paul S, Kumar P, et al. (2018) Diagnostic utility of bone marrow aspiration, imprint and biopsy in evaluation of various haematological disorders. Int J Curr Res Med Sci 4: 28-36.
- Vijayamohanan L, Asotra S, Kumari K, Murgai P, Dattal D (2020) Comparative utility of bone marrow aspiration and trephine biopsy in evaluation of hematological disorders. Arch Med Health Sci 8: 15-19.
- Wilkins BS (2011) Lymphoma. In: Porwit, Mc Cullough J, Erber WN, Blood and bone marrow pathology. (2nd edn), Edinburgh: Elsevier Limited, 412-442.
- Zhang QY, Foucar K (2009) Bone marrow involvement by Hodgkin and non-Hodgkin lymphomas. Hematol Oncol Clin North Am 23: 873-902.
- Graf LB, Korte W, Schmid L, Schmid U, Cogliatti SB (2005) Impact of aspirate smears and trephine biopsies in routine bone marrow diagnostics: A comparative study of 141 cases. Swiss Med Wkly 135: 151-159.
- Chauhan S, Pradhan S, Mohanty R, Saini A, Devi K, et al. (2018) Evaluation of sensitivity and specificity of bone marrow trephine biopsy tests in an Indian teaching hospital. Alex J Med 54: 161-166.