

Review Article

THE USE OF POST-OPERATIVE ANALGESICS IN ORTHOPEDIC PRACTICES: A COMPREHENSIVE REVIEW

Zulfiqar Haider Kerawala¹

¹Specialist A Orthopedic Surgeon, Al Qassimi Hospital, Sharjah, United Arab Emirates.

Received : 02/09/2025
Received in revised form : 06/10/2025
Accepted : 22/10/2025

Corresponding Author:

Dr. Zulfiqar Haider Kerawala,
Consultant Orthopaedic Surgeon,
Sandeman Provincial Hospital, Quetta
Pakistan.
Email: zhkerawala@gmail.com

DOI: 10.70034/ijmedph.2025.4.206

Source of Support: Nil,
Conflict of Interest: None declared

Int J Med Pub Health
2025; 15 (4); 1150-1155

ABSTRACT

Pain management after surgery is a key component of orthopedic practice which eventually has a direct impact on surgical results, recovery, and patient satisfaction. In this comprehensive review, the use of post-operative analgesics in orthopedic surgery is defined. It highlights the importance of managing pain after surgery using both, non-pharmacological and pharmacological approaches. The function of multimodal analgesia is discussed in this review which reduces the dependence of opioid and associated negative effects. It also improves pain management by merging several regional anesthetic procedures and analgesic drugs. Patient-specific factors are also covered in this review which include pain management plans according to age, medication sensitivity, and comorbidities. It also looks over advanced developments such as digital pain monitoring devices, sustained-release medication formulations, innovative analgesic compounds, and patient-centered treatment methods.

Keywords: Pain, postoperative, nonpharmacological, pharmacological.

INTRODUCTION

Orthopedic surgery is a unique sub-category of medicine which is involved in the treatment of trauma or conditions related to muscle tissues, bone, and joints.^[1] It includes a wide range of activities such as fracture management, and joint replacements. After orthopedic surgery, the first thing that must to be taken care of is the management of pain the patient is experiencing.^[2] The most common feeling after an individual's orthopedic surgery is pain as this treatment includes the operation of bones, soft tissues, and joints. If not managed properly, this pain leads to a slower recovery, develops a long-term chronic pain, and eventually affects the quality of life of the individual. Pain after orthopedic surgery can be complicated.^[3] This pain is influenced by a variety of factors such as the amount of tissue which was damaged or manipulated, surgery technique, and the individual's particular circumstances. Patients might go through muscle spasms, sharp acute pain, and swelling (inflammation). Consequently, the provision of efficient pain relief, which is also referred to as analgesia, becomes the most important aspect of the patient's care. The failure to manage pain effectively could result in delays in patients

being able to move about and recover, a decrease in their physical function, and a longer stay in the hospital.

The impact of efficient pain management after orthopedic surgery on patient recovery and satisfaction is quite significant. Adequate analgesia during the postoperative period not only comforts patients but also allows them to get involved in active therapies, thus facilitating normal daily activities quicker.^[4] Good pain management practices can also help to minimize complications, and speed up the healing process. Therefore, it has become an indispensable component of orthopedic patient care. Conversely, if pain is not properly managed, it will certainly lead to increased healthcare costs.^[5] Patients suffering from pain that cannot be controlled may need longer hospital stays, more treatments, and greater or different medication. Therefore, poor pain control is not only a medical issue but also a financial concern for hospitals and the healthcare systems. This review presents the various methods of pain management following surgery that are currently employed in the orthopedic practice. Its purpose is to illustrate both drug (pharmacological) and non-drug (non-pharmacological) methods of controlling postoperative pain.^[6] Moreover, the review will explore the latest advancements and breakthroughs in

this area. It will feature the major orthopedic surgeries that often necessitate pain management, the objectives of pain control, and the most common medications. In addition to the classic opioid and non-opioid medications, the review will explore regional anesthesia techniques like epidural analgesia and nerve blocks, as well as the concept of multimodal analgesia — a strategy that combines different pain relief measures for optimal outcomes.

REVIEW

Goals of postoperative analgesia in orthopedic practice

Objectives of Pain Management

Relieving the patient's pain totally is the primary objective of postoperative pain treatment. Reducing pain to a manageable and pleasant level is the main goal. Patients can walk, breathe, and participate in rehabilitation with less discomfort when their pain is adequately managed. Instead of battling chronic pain, patients who are not in excruciating agony can concentrate their energies on recovering and eating healthily.^[7] A key component of rehabilitation following surgery is early mobility, which is made feasible by efficient pain management. Patients in significant pain can be hesitant or fearful to move, which can lead to significant consequences including pressure ulcers (bed sores), deep vein thrombosis (blood clots), or atelectasis (lung collapse). Patients who receive effective pain management interventions will move sooner and more frequently. Early mobility maintains healthy lung function, vascular circulation, and minimizes postoperative complications.^[8]

Pain control is an important part of care for surgical patients. It will impact the comfort of the patient, but can also have an impact on the ultimate outcome for the patient. Patients who are pain-free will be more likely to follow the postoperative course prescribed for them, which may include rehabilitation exercises or physical therapy.^[9] An active patient will heal more quickly, have more mobility, and have improved overall results in the long-term. Decreasing pain is only one of two components associated with appropriate pain management. The other component is to prevent or minimize unwanted side effects from the analgesics used to treat pain. Constipation is one unwanted side effect of opioids, and respiratory depression (or decreased respiratory rate) is another. Finding the very best situation in which pain control and side effects are tightly controlled is significant.^[10] Patient satisfaction is a key part of postoperative care. A speedy and successful recovery can significantly impact a patient's perception of their care, as well as their overall happiness with the hospital or clinic. Excellent patient satisfaction hinges on efficient pain management. When patients are comfortable and have adequate pain management after an operation, they are more likely to have a positive perception of

their care, enhancing recovery as well as the positive reputation of the hospital or clinic.^[11]

Factors that Influence the Choice of Analgesic Agents

The type of orthopedic surgery will guide whether analgesics are needed. A comprehensive and aggressive strategy of pain management is used in major joint replacements such as knee or hip replacements. These often utilize regional anesthetic procedures with opioids. On the other hand, mild analgesics may suffice in smaller procedures such as arthroscopic procedures. These mild analgesics include a non-steroidal anti-inflammatory which is called NSAID. This reduces the risk of side effects that are linked with opioids and provide adequate pain relief. An individual's specific needs should also be considered as part of the pain management plan.^[12] This includes their age as older patients are prone to the negative effects of opioids. Additionally, patients with other medical problems, like heart or lung issues, which may stay stable, will also be cautious about the medication selection to prevent worsening ongoing symptoms. Also, it's important to remember to look for drug sensitivities or allergies to the medications chosen. If the possible medication is not tolerated, then more alternative methods are to be employed for pain management, in order to avoid negative side effects and ensure patient safety.^[13]

The level of pain that an individual reports after a surgery is also considered in selecting the pain medication. The more the pain would be, the stronger medicine will be appropriate for it. Mostly, NSAIDs and acetaminophen are sufficient for managing mild pain as they are considered non-opioid medications. They are usually linked with minor and less invasive orthopedic procedures. Where there is severe pain, opioids are used. They are the standard medicines after surgical procedures. Other than this, side effects must be considered. It is important to weigh the pain relief and side effects to ensure the correct selection of medicine.^[14]

Balanced Approach to Analgesia

It is important to manage pain in a personalized manner because every individual experiences pain differently. Every patient's need is different. Therefore, while selecting the medication to relieve pain, it is important to consider the type, dose, and method of delivering pain medication. Moreover, it is important to ensure that the side effects are reduced and pain is controlled. A combination of techniques and medicines is called multimodal analgesia. This helps in managing the pain through different techniques, ultimately reducing the dependence on opioids. Furthermore, it is important to adjust the doses based on the patient's response to provide them effective and safe relief. This can be tracked through continuous monitoring and proper titration. Patients should also get educated about their possible side effects, and pain management options. This enhances communication and empowers the patients to take an active role in their recovery. Lastly, if pain is managed and treated early and on time, it supports

faster healing, prevents the pain from worsening, and improves comfort.^[15]

Pharmacological analgesic agents

Non-opioid Analgesics

NSAIDs (non-steroidal anti-inflammatory drugs) work by inhibiting the work of the cyclooxygenase (COX) enzymes, which function to produce fever, pain, and inflammation-causing chemicals called prostaglandins, from arachidonic acid. NSAIDs relieve pain and inflammation by inhibiting prostaglandin production. In the orthopedic management of pain, NSAIDs are frequently used for mild to moderate postoperative pain relief and inflammation. NSAIDs are especially useful for pain conditions such as osteoarthritis, tendinitis, and following surgeries in which inflammation is the primary mechanism causing pain. NSAIDs are commonly used as part of multimodal analgesia, which is to integrate pain medications to reduce the dependence on opioids. This serves two purposes: reducing the side effects caused by dependence on opioids and integrating a variety of medications to manage pain symptoms. Patients enjoy a more comfortable and satisfying recovery with this balance.^[16]

Acetaminophen, or paracetamol, mainly affects the brain. By inhibiting the central nervous system from producing prostaglandins via the hypothalamus, it lowers fever and pain. Acetaminophen is the most commonly used and preferred first-line treatment for mild to severe pain. It is particularly useful for patients that are unable to take NSAIDs due to gastrointestinal issues (ulcers, bleeding risk). Acetaminophen is tolerated better and has lower gastrointestinal adverse effects compared to NSAIDs for individuals with these problems. In orthopedic postoperative therapy, it is routinely used orally and can be used as a part of a multimodal pain plan to reduce opioid use and provide adequate pain control.^[17]

Opioid Analgesics

Morphine is an effective opioid analgesic that works by binding to μ -opioid receptors in the spinal cord and brain. This changes the perception of pain in the body along with sedation and relief of pain. In orthopedic care, mostly morphine is used to manage severe pain after surgery. That pain does not respond to non-opioids such as NSAIDs or acetaminophen. Just like morphine, there is another semi-synthetic analgesic called oxycodone which binds to μ -opioid receptors in the central nervous system in order to decrease the pain. Oxycodone is mostly used to manage moderate to severe pain. Oxycodone is included in a multimodal approach to reduce pain. It is included as a part of a regimen with other analgesics. By using this multimodal approach, the amount of opioids needed is reduced, pain relief is improved, and side effects such as constipation and sedation are prevented. Hydrocodone is another semi-synthetic opioid which also reduces pain by binding to μ -opioid receptors. To treat acute pain after surgery, especially when non-opioid analgesics

are inadequate, hydrocodone is used. It is mostly used with NSAIDs or acetaminophen to decrease pain as well as the use of strong opioids.^[18]

Adjuvant Analgesics

The use of anticonvulsant drugs like gabapentin and pregabalin is very successful in the management of neuropathic pain which is the pain that occurs due to nerve irritation or injury. The mechanism of these medications is by calcium channels in the brain and spinal cord alterations leading to the reduction of neurotransmitters such as glutamate that are pain signalers. These drugs are commonly prescribed after the orthopedic procedure as adjunct therapy particularly when nerve pain emerges as a result of compression or injury during surgery [19]. The role of these drugs in pain therapy is that they are effective in treating the pain caused by nerve damage which may not be responding to conventional pain medications.

Tricyclic Antidepressants

Tricyclic antidepressants (TCAs) like amitriptyline, may be utilized for the treatment of other pains as well. The ultimate goal of the drugs is to facilitate the process of the two neurotransmitters, serotonin and norepinephrine, by blocking their reabsorption in the brain. The increased concentration of the two neurotransmitters in the brain, along with the reduced pain perception, is the ultimate goal of using such medications. Occasionally, as with anticonvulsants, these categories of antidepressants are employed for the treatment of neurologic or chronic pain occurring after orthopedic surgeries, especially when nerve injury or damage is implicated. The use of TCAs can be in the facilitation of both comfort and long-term outcomes, as this type of pain is often very hard to manage with ordinary pain relievers.^[20]

Anticonvulsants

By regulating neuronal activity in the brain and spinal cord, many anticonvulsant medications, such as lamotrigine and carbamazepine, can also help reduce pain. They lessen the aberrant nerve impulses that result in neuropathic pain. Although they are not as commonly used as other painkillers, these medications may be useful if nerve pain contributes significantly to postoperative discomfort, particularly if standard painkillers such as acetaminophen, opioids, or NSAIDs are insufficient.^[21]

Regional anesthesia techniques

Epidural Analgesia

An extremely successful pain management technique is epidural analgesia. In order to specifically relieve lower body pain, local anesthetics are injected into the epidural region around the spinal cord. This method is frequently applied to postoperative pain control and childbirth. The early mobilization and increased comfort for the patient resulting from appropriate epidural analgesia promote quicker recovery. One of the main advantages is a reduction in systemic opioid medications, therefore minimizing adverse effects such as lethargy, nausea, and constipation.^[22]

Peripheral Nerve Blocks

Peripheral nerve blocks are a method to alleviate pain in a specific area by placing anesthetics very close to the nerves that transmit pain signals. This method causes loss of sensation in a particular area such as the arm or leg while the rest of the body is intact. Nerve blocks can be used to eliminate pain signals effectively and, at the same time, decrease the demand for opioids considerably along with the risks associated with opioid use, i.e., respiratory problems and addiction. Patients in recovery generally feel more awake and display fewer side effects. Properly managed regional pain control is one of the factors that contribute to the patient's mobility and hence, the early rehabilitation, which in turn, reduces the incidence of complications like deep vein thrombosis and muscle atrophy. Additionally, it promotes patients' general comfort and motivates them to take part in their recovery process more.^[23]

Intrathecal Analgesia

During intrathecal analgesia, medication is injected straight into the cerebrospinal fluid that is in the spinal canal. It is a good option for postoperative care as it offers prompt and potent pain relief. Furthermore, the incidence of systemic side effects such as nausea, constipation, and respiratory depression is markedly less as the medication acts on the spinal nerves only. Moreover, this technique enables doctors to closely monitor the dosage, thus providing pain relief that is both trustworthy and safe. This selective technique not only enhances patient comfort but also facilitates quicker recovery after surgery without the patient being aware of the pain.^[24]

Multimodal analgesia

Rationale for Multimodal Analgesia

Pain is caused by several pathways and mechanisms. Multimodal analgesia, which uses different analgesic drugs targeting different parts of the pain pathway, solves this problem. This technique promises more extensive and efficient pain management than single medication use. It is particularly beneficial in orthopedic practice since patients commonly suffer from both neuropathic and nociceptive pain. It limits opioid use, thereby diminishing the chances of addiction and the coming along issues related to opioids. This method makes it possible to have safe and more extensive pain management by mixing local techniques with non-opioid medications. As treatment plans can be customized to fit each patient's needs, preferences, and medical conditions, this method also offers personalized care.^[25] Personalized multimodal therapy leads to better recovery outcomes, patient comfort, and satisfaction.

Combination Therapy Using Various Analgesic Agents

Acetaminophen and NSAIDs are examples of non-opioid painkillers that are used in multimodal analgesia. These drugs help control mild-to-moderate pain and inflammation while lowering the need for opioids. Opioids like morphine, oxycodone, and hydrocodone are only used for moderate-to-severe pain when combined with other medications. This

reduces the risk of respiratory depression, constipation, and exhaustion while providing effective pain relief. Targeted pain control for certain surgical sites is made possible by regional anesthesia procedures such as epidural, peripheral nerve blocks, and intrathecal analgesia.^[26] These techniques diminish the need for systemic drugs, promote early mobility, and minimize side effects. For the treatment of neuropathic pain following orthopedic procedures, adjuvant drugs such as pregabalin, gabapentin, tricyclic antidepressants, and anticonvulsants are especially helpful.

Benefits and Challenges of Multimodal Analgesia

There are a few benefits of multimodal analgesia. First of all, it works through several methods to improve pain relief. Due to this, patients are mostly satisfied, recover quickly, and have greater comfort. It reduces the need of high-dose opioids which encourages safer post-operative care and reduces side effects. In orthopedic recovery, rehabilitation and early mobilization are supported by better pain management.^[27]

Along with some benefits, there are a few challenges as well that are linked with multimodal analgesia. To implement the multimodal analgesia, it requires accurate dosing, careful planning, and continuous monitoring. Moreover, personalized evaluation is important as every patient responds differently to medications. Multimodal approaches are also costly initially.

Patient-specific considerations

Age, Gender, and Pain Perception

Age is a major factor which influences an individual's reaction to analgesics and perception of pain. As age increases, metabolism becomes slow. Therefore, older people mostly need low doses of medicines to prevent adverse effects such as opioid-induced respiratory depression. In order to guarantee efficacy and safety, patients require doses according to their age. Gender also affects how people react to medicine and perceive pain. According to other research, females are more sensitive to pain which influences the dose and selection of analgesics.^[28]

Coexisting Medical Conditions

Planning for pain treatment requires taking preexisting medical issues into account. Certain NSAIDs should be avoided by patients with cardiovascular problems since they increase the risk of heart attacks and strokes. Lower dosages or non-opioid alternatives may be more beneficial for people with respiratory conditions, such as COPD, who are more susceptible to breathing issues caused by opioids. Dosage modifications are necessary for patients with liver or kidney impairment in order to avoid medication toxicity and buildup. Patients with diabetes may experience pain differently and heal wounds more slowly, necessitating close blood sugar monitoring and pain management to promote healing.^[29]

Medication Allergies and Sensitivities

Prior to beginning treatment, it is essential to identify drug sensitivities. To prevent responses ranging from minor rashes to severe anaphylaxis, patients with known allergies should be given suitable substitutes. Even if they do not have allergies, some people may be sensitive to or intolerant of particular analgesics. Alternative agents or techniques should be employed in these situations, backed by careful observation and patient communication.^[30]

Emerging trends and innovations

New Analgesic Agents and Delivery Methods

New analgesic drugs with sophisticated modes of action, like ion channel blockers, NMDA receptor antagonists, and selective opioid receptor modulators, have been developed recently. Due to this, pain is managed effectively with very little adverse effects. This ultimately benefits the recovery of the patient after surgery. Developments in drug delivery systems and sustained-release formulations provide less doses and long-lasting pain relief. In order to treat localized pain, topical analgesics are being used in a big quantity. Topical analgesics include gels, creams, and patches. These are especially used for musculoskeletal disorders such as osteoarthritis. These analgesics reduce systemic side effects and focus on providing tailored pain treatment where needed.^[31]

Advances in Pain Assessment and Monitoring

Due to recent digital health technologies such as smartphone applications and other wearables, it has become possible to track real-time pain. Individuals can record their pain levels which will help the healthcare workers to customize their treatment plans based on the ongoing data. Some objective pain measures are introduced, that include imaging methods and biomarkers, which helps in determining the accurate severity and location of pain. These techniques produce accurate statistics and reduce the reliance on self-reporting.^[32] Due to telemedicine, pain management experts are now more accessible specially for patients who are in remote areas. It promotes easy follow-ups, expert-guided care, and increases the participation of the patient.

Patient-Centered Care in Pain Management

Healthcare professionals and individuals have to work together to make better decisions regarding contemporary pain treatment. This increases involvement in care and confidence because the treatment programs are designed according to the patient's requirements, preferences, and objectives. Patient education is the foundation of this strategy. Patients are allowed to take an active role in their recovery. If the patients are aware of the negative consequences of the options available to them and their after effects, they can make educated decisions. The main goal of this patient-centered care is to manage pain effectively and have a quality life. The purpose is to restore mental well-being, comfort, and function along with pain relief. These plans are customized based on each patient's lifestyle, and health status.

CONCLUSION

After orthopedic surgery, it is important to manage the pain effectively in order to achieve better results and guarantee a satisfying surgical experience. This study emphasized on the main facets of pain management in orthopedic surgery. It concludes that if pain is managed properly, it leads to early movement, accelerates recovery, and increases patient comfort. Multimodal analgesia offers a well-rounded approach to manage the pain which blends a few methods and painkillers. Some of the recent advancements in the management of pain include sophisticated pain assessment instruments, novel analgesic drugs, and patient-centered case methodologies. These advancements make the treatment efficient and safer.

REFERENCES

- Goel S, Deshpande SV, Jadawala VH, Suneja A, Singh R. A comprehensive review of postoperative analgesics used in orthopedic practice. *Cureus*. 2023 Nov 13;15(11).
- Aldanyowi SN. Novel techniques for musculoskeletal pain management after orthopedic surgical procedures: a systematic review. *Life*. 2023 Dec 15;13(12):2351.
- Chunduri A, Aggarwal AK. Multimodal pain management in orthopedic surgery. *Journal of clinical medicine*. 2022 Oct 28;11(21):6386.
- Argoff CE. Recent management advances in acute postoperative pain. *Pain Practice*. 2014 Jun;14(5):477-87.
- Chou R, Gordon DB, de Leon-Casasola OA, Rosenberg JM, Bickler S, Brennan T, Carter T, Cassidy CL, Chittenden EH, Degenhardt E, Griffith S. Management of Postoperative Pain: a clinical practice guideline from the American pain society, the American Society of Regional Anesthesia and Pain Medicine, and the American Society of Anesthesiologists' committee on regional anesthesia, executive committee, and administrative council. *The journal of pain*. 2016 Feb 1;17(2):131-57.
- Roberts M, Brodribb W, Mitchell G. Reducing the pain: a systematic review of postdischarge analgesia following elective orthopedic surgery. *Pain Medicine*. 2012 May 1;13(5):711-27.
- Ojha A, Xia R, Wei X, Karn R, Tamrakar S. Post-operative pain management.
- Sampognaro G, Harrell R. Multimodal postoperative pain control after orthopaedic surgery.
- Strassels SA, McNicol E, Suleman R. Postoperative pain management: a practical review, part 2. *American Journal of Health-System Pharmacy*. 2005 Oct 1;62(19):2019-25.
- Strassels SA, McNicol E, Suleman R. Postoperative pain management: a practical review, part 1. *American Journal of Health-System Pharmacy*. 2005 Sep 15;62(18):1904-16.
- Gupta A, Kaur K, Sharma S, Goyal S, Arora S, Murthy RS: Clinical aspects of acute post-operative pain management & its assessment. *J Adv Pharm Technol Res*. 2010, 1:97-108.
- Abebe MM, Arefayne NR, Temesgen MM, Admass BA: Evidence-based perioperative pain management protocol for day case surgery in a resource limited setting: systematic review. *Ann Med Surg (Lond)*. 2022, 80:104322. 10.1016/j.amsu.2022.104322
- Queremel Milani DA, Davis DD: Pain Management Medications. StatPearls Publishing, Treasure Island, FL; 2023.
- Buli B, Gashaw A, Gebeyehu G, Abrar M, Gerbessa B: Patient satisfaction with post-operative pain management and associated factors among surgical patients at Tikur Anbessa Specialized Hospital: crosssectional study. *Ann Med Surg (Lond)*. 2022, 79:104087. 10.1016/j.amsu.2022.104087
- Anastase DM, Cionac Florescu S, Munteanu AM, Ursu T, Stoica CI: Analgesic techniques in hip and knee arthroplasty:

- from the daily practice to evidence-based medicine. *Anesthesiol Res Pract.* 2014, 2014:569319. 10.1155/2014/569319
16. Chau DL, Walker V, Pai L, Cho LM: Opiates and elderly: use and side effects. *Clin Interv Aging.* 2008, 3:273- 8. 10.2147/cia.s1847
 17. Anekar AA, Hendrix JM, Cascella M: WHO Analgesic Ladder. StatPearls Publishing, Treasure Island, FL; 2023.
 18. Sugai DY, Deptula PL, Parsa AA, Don Parsa F: The importance of communication in the management of postoperative pain. *Hawaii J Med Public Health.* 2013, 72:180-4.
 19. Wells N, Pasero C, McCaffery M: Improving the quality of care through pain assessment and management. *Patient Safety and Quality: An Evidence-Based Handbook for Nurses.* Hughes RG (ed); Agency for Healthcare Research and Quality, Rockville, MD; 2008.
 20. Gunaydin C, Bilge SS: Effects of nonsteroidal anti-inflammatory drugs at the molecular level. *Eurasian J Med.* 2018, 50:116-21. 10.5152/eurasianjmed.2018.0010
 21. Ayoub SS: Paracetamol (acetaminophen): a familiar drug with an unexplained mechanism of action. *Temperature (Austin).* 2021, 8:351-71. 10.1080/23328940.2021.1886392
 22. Pathan H, Williams J: Basic opioid pharmacology: an update. *Br J Pain.* 2012, 6:11-6. 10.1177/2049463712438493
 23. Rogawski MA, Löscher W, Rho JM: Mechanisms of action of antiseizure drugs and the ketogenic diet. *Cold Spring Harb Perspect Med.* 2016, 6:a022780. 10.1101/cshperspect.a022780
 24. Olawin AM, M Das J: Spinal Anesthesia. StatPearls Publishing, Treasure Island, FL; 2023.
 25. Karcz M, Papadakos PJ: Respiratory complications in the postanesthesia care unit: a review of pathophysiological mechanisms. *Can J Respir Ther.* 2013, 49:21-9.
 26. Chang A, Dua A, Singh K, White BA: Peripheral Nerve Blocks. StatPearls Publishing, Treasure Island, FL; 2023.
 27. Pehora C, Pearson AM, Kaushal A, Crawford MW, Johnston B: Dexamethasone as an adjuvant to peripheral nerve block. *Cochrane Database Syst Rev.* 2017, 2017:CD011770. 10.1002/14651858.CD011770.pub2
 28. Manworren RC: Multimodal pain management and the future of a personalized medicine approach to pain. *AORN J.* 2015, 101:308-14; quiz 315-8. 10.1016/j.aorn.2014.12.009
 29. Folino TB, Mahboobi SK: Regional Anesthetic Blocks. StatPearls Publishing, Treasure Island, FL; 2023.
 30. Knotkova H, Pappagallo M: Adjuvant analgesics. *Med Clin North Am.* 2007, 91:113-24. 10.1016/j.mcna.2006.10.007
 31. Bhatia A, Buvanendran A: Anesthesia and postoperative pain control-multimodal anesthesia protocol. *J Spine Surg.* 2019, 5:S160-5. 10.21037/jss.2019.09.33
 32. Fenske JN, Berland DW, Chandran S, et al.: Pain Management. Michigan Medicine University of Michigan, Ann Arbor, MI; 2021.